arXiv:1401.1108v2 [math.NT] 8 Jan 2014

Proof of a conjecture related to divisibility properties of binomial coefficients*

Quan-Hui Yang†

School of Mathematics and Statistics,

Nanjing University of Information Science and Technology,

Nanjing 210044, P. R. China

Abstract

Let a, b and n be positive integers with $a > b$. In this note, we prove that

$$(2bn+1)(2bn+3)\binom{2bn}{bn}\Big|3(a-b)(3a-b)\binom{2an}{an}\binom{an}{bn}.$$

This confirms a recent conjecture of Amdeberhan and Moll.

2010 Mathematics Subject Classifications: 11B65, 05A10

Keywords: binomial coefficients, p-adic order, divisibility properties

1 Introduction

Let $\mathbb{Z}$ denote the set of all integers. In 2009, Bober [1] determined all cases such that

$$\frac{(a_1n)!\cdots(a_kn)!}{(b_1n)!\cdots(b_{k+1}n)!}\in\mathbb{Z},$$

*This work was supported by the National Natural Science Foundation of China, Grant No. 11371195.

†Email: yangquanhui01@163.com.

where $a_s \neq b_t$ for all s, t, $\sum a_s = \sum b_t$ and $\gcd(a_1, \ldots, a_k, b_1, \ldots, b_{k+1}) = 1$.

Recently, Z.-W. Sun [12, 13] studied divisibility properties of binomial coefficients and obtained some interesting results. For example,

$$2(2n+1)\binom{2n}{n} \Big| \binom{6n}{3n}\binom{3n}{n},$$

$$(10n+1)\binom{3n}{n} \Big| \binom{15n}{5n}\binom{5n-1}{n-1}.$$

Later, Guo and Krattenthaler (see [5, 7]) got some similar divisibility results. For related results, one can refer to [2]-[4] and [8]-[11].

Let

$$S_n = \frac{\binom{6n}{3n}\binom{3n}{n}}{2(2n+1)\binom{2n}{n}} \quad \text{and} \quad t_n = \frac{\binom{15n}{5n}\binom{5n-1}{n-1}}{(10n+1)\binom{3n}{n}}.$$

In [6], Guo proved the following Sun's conjectures.

Theorem A. (See [12, Conjecture 3(i)].) *Let n be a positive integer. Then*

$$3S_n \equiv 0 \pmod{2n+3}.$$

Theorem B. (See [13, Conjecture 1.3].) *Let n be a positive integer. Then*

$$21t_n \equiv 0 \pmod{10n+3}.$$

Some other results are also proved.

T. Amdeberhan and V. H. Moll proposed the following conjecture (See Guo [6, Conjecture 7.1]).

Conjecture 1. *Let a, b and n be positive integers with $a > b$. Then*

$$(2bn+1)(2bn+3)\binom{2bn}{bn} \Big| 3(a-b)(3a-b)\binom{2an}{an}\binom{an}{bn}.$$

In this note, we give the proof of this conjecture.

Theorem 1. *Conjecture 1 is true.*

Remark 1. *Let $a = 3$ and $b = 1$. Then Theorem A follows from Theorem 1 immediately.*

2 Proofs

For an integer n and a prime p, we write $p^k \| n$ if $p^k | n$ and $p^{k+1} \nmid n$. We use $\nu_p(n)$ to denote such integer k. It is well known that

$$\nu_p(n!) = \sum_{i=1}^{\infty} \left\lfloor \frac{n}{p^i} \right\rfloor, \tag{1}$$

where $\lfloor x \rfloor$ denotes the greatest integer not exceeding x.

Before the proof of Theorem 1, we give the following lemma.

Lemma 1. *Let x and y be two real numbers. Then $\lfloor 2x \rfloor + \lfloor y \rfloor \geq \lfloor x \rfloor + \lfloor x - y \rfloor + \lfloor 2y \rfloor$.*

Proof. Noting that $2x + y = x + (x - y) + 2y$, we only need to prove that $\{2x\} + \{y\} \leq \{x\} + \{x - y\} + \{2y\}$, where $\{z\}$ denotes the fractional part of z. We can prove it by comparing $\{x\}$ and $\{y\}$ with $1/2$. We leave the proof to the reader. □

Proof of Theorem 1. Let

$$T(a, b, n) := \binom{2an}{an}\binom{an}{bn} \Big/ \binom{2bn}{bn} = \frac{(2an)!(bn)!}{(an)!(an - bn)!(2bn)!}.$$

By (1), for any prime p, we have

$$\nu_p(T(a, b, n)) = \sum_{i=1}^{\infty} \left(\left\lfloor \frac{2an}{p^i} \right\rfloor + \left\lfloor \frac{bn}{p^i} \right\rfloor - \left\lfloor \frac{an}{p^i} \right\rfloor - \left\lfloor \frac{an - bn}{p^i} \right\rfloor - \left\lfloor \frac{2bn}{p^i} \right\rfloor \right).$$

By Lemma 1, it follows that each term of $\nu_p(T(a, b, n))$ is nonnegative. Hence $\nu_p(T(a, b, n)) \geq 0$. Therefore, $T(a, b, n) \in \mathbb{Z}$.

Since $\gcd(2bn + 1, 2bn + 3) = 1$, it suffices to prove that

$$2bn + 1 | 3(a - b)(3a - b)T(a, b, n)$$

and

$$2bn + 3 | 3(a - b)(3a - b)T(a, b, n).$$

Now we first prove the latter part. Suppose that $p^\alpha \| 2bn+3$ with $\alpha \geq 1$. Then we shall prove

$$p^\alpha | 3(a-b)(3a-b)T(a,b,n). \tag{2}$$

Let $p^\beta \| a-b$ and $p^\gamma \| 3a-b$ with $\beta \geq 0$ and $\gamma \geq 0$. Write $\tau = \max\{\beta, \gamma\}$. If $\alpha \leq \tau$, then (2) clearly holds. Now we assume $\alpha > \tau$.

Suppose that $p \geq 5$. Next we prove

$$\left\lfloor \frac{2an}{p^i} \right\rfloor + \left\lfloor \frac{bn}{p^i} \right\rfloor - \left\lfloor \frac{an}{p^i} \right\rfloor - \left\lfloor \frac{an-bn}{p^i} \right\rfloor - \left\lfloor \frac{2bn}{p^i} \right\rfloor = 1$$

for $i = \tau+1, \tau+2, \ldots, \alpha$. Noting that $p|2bn+3$ and $p \geq 5$, we have $\gcd(p,n) = 1$. Otherwise, we have $p|3$, a contradiction.

By $p^\alpha \| 2bn+3$, it follows that $2bn \equiv p^\alpha - 3 \pmod{p^\alpha}$ and $bn \equiv (p^\alpha - 3)/2 \pmod{p^\alpha}$.

Take an integer $i \in \{\tau+1, \tau+2, \ldots, \alpha\}$. Then $2bn \equiv p^i - 3 \pmod{p^i}$ and $bn \equiv (p^i-3)/2 \pmod{p^i}$. Now we divide into several cases according to the value of $an \pmod{p^i}$.

Case 1. $an \equiv t \pmod{p^i}$ with $0 \leq t < (p^i-3)/2$. It follows that $2an \equiv 2t \pmod{p^i}$ and $0 \leq 2t < p^i - 3$. We also have

$$an - bn \equiv t - (p^i-3)/2 + p^i \pmod{p^i},$$

where $0 \leq t - (p^i-3)/2 + p^i < p^i$. Hence

$$\begin{aligned} & \left\lfloor \frac{2an}{p^i} \right\rfloor + \left\lfloor \frac{bn}{p^i} \right\rfloor - \left\lfloor \frac{an}{p^i} \right\rfloor - \left\lfloor \frac{an-bn}{p^i} \right\rfloor - \left\lfloor \frac{2bn}{p^i} \right\rfloor \\ = \; & \frac{2an-2t}{p^i} + \frac{bn-(p^i-3)/2}{p^i} - \frac{an-t}{p^i} \\ & - \left(\frac{an-bn-(t-(p^i-3)/2+p^i)}{p^i} \right) - \frac{2bn-(p^i-3)}{p^i} \\ = \; & 1. \end{aligned}$$

Case 2. $an \equiv (p^i-3)/2 \pmod{p^i}$. Clearly, $an - bn \equiv 0 \pmod{p^i}$. Since $\gcd(p,n) = 1$, we have $p^i | a-b$. However, $p^\beta \| a-b$ and $\beta \leq \tau < i$, a contradiction.

Case 3. $an \equiv (p^i-1)/2 \pmod{p^i}$. It follows that

$$3an - bn \equiv \frac{3(p^i-1)}{2} - \frac{(p^i-3)}{2} \equiv 0 \pmod{p^i}.$$

By $\gcd(p,n)=1$, we have $p^i | 3a-b$. This contradicts $p^\gamma \| 3a-b$ and $\gamma < i$.

Case 4. $an \equiv t \pmod{p^i}$ with $(p^i+1)/2 \le t < p^i$. It follows that

$$2an \equiv 2t - p^i \pmod{p^i}, \quad 0 \le 2t - p^i < p^i.$$

We also have

$$an - bn \equiv t - (p^i-3)/2 \pmod{p^i}, \quad 0 \le t-(p^i-3)/2 < p^i.$$

Hence

$$\begin{aligned} & \left\lfloor \frac{2an}{p^i} \right\rfloor + \left\lfloor \frac{bn}{p^i} \right\rfloor - \left\lfloor \frac{an}{p^i} \right\rfloor - \left\lfloor \frac{an-bn}{p^i} \right\rfloor - \left\lfloor \frac{2bn}{p^i} \right\rfloor \\ = & \frac{2an-(2t-p^i)}{p^i} + \frac{bn-(p^i-3)/2}{p^i} - \frac{an-t}{p^i} \\ & - \left(\frac{an-bn-(t-(p^i-3)/2)}{p^i} \right) - \frac{2bn-(p^i-3)}{p^i} \\ = & 1. \end{aligned}$$

Therefore, $\nu_p(T(a,b,n)) \ge \alpha - \tau$, and then

$$\nu_p(3(a-b)(3a-b)T(a,b,n)) \ge \alpha.$$

That is, (2) holds.

Now we deal with the case $p=3$. If $9|n$, then $3|2bn+3$ and $9 \nmid 2bn+3$. It follows that $\alpha = 1$, and then (2) clearly holds. If $9 \nmid n$, then we can follow the proof of the case $p \ge 5$ above. In Case 2, by $an - bn \equiv 0 \pmod{3^i}$, we have $3^{i-1} | a-b$. In Case 3, we have $3^{i-1} | 3a-b$. So, if $i \ge \tau+2$, then $i-1 \ge \tau+1$. It is a contradiction in both cases. Hence

$$\left\lfloor \frac{2an}{3^i} \right\rfloor + \left\lfloor \frac{bn}{3^i} \right\rfloor - \left\lfloor \frac{an}{3^i} \right\rfloor - \left\lfloor \frac{an-bn}{3^i} \right\rfloor - \left\lfloor \frac{2bn}{3^i} \right\rfloor = 1$$

for $i = \tau+2, \tau+3, \ldots, \alpha$. It follows that $\nu_3(T(a,b,n)) \ge \alpha - \tau - 1$, and then

$$\nu_3(3(a-b)(3a-b)T(a,b,n)) \ge \alpha.$$

That is, (2) also holds. Hence, $2bn+3|3(a-b)(3a-b)T(a,b,n)$.

The proof of $2bn+1|3(a-b)(3a-b)T(a,b,n)$ is very similar. We omit it here.

This completes the proof of Theorem 1.

□

References

[1] J.W. Bober, *Factorial ratios, hypergeometric series, and a family of step functions,* J. Lond. Math. Soc. 79 (2009), 422-444.

[2] N.J. Calkin, *Factors of sums of powers of binomial coefficients,* Acta Arith. 86 (1998), 17-26.

[3] H.Q. Cao and H. Pan, *Factors of alternating binomial sums,* Adv. in Appl. Math. 45 (2010), 96-107.

[4] N. Fine, *Binomial coefficients modulo a prime,* Amer. Math. Monthly 54 (1947), 589-592.

[5] V.J.W. Guo, *Proof of Suns conjecture on the divisibility of certain binomial sums,* Electron. J. Combin. 20(4) (2013), #P20.

[6] V.J.W. Guo, *Proof of two divisibility properties of binomial coefficients conjectured by Z.-W. Sun,* available at http://arxiv.org/abs/1312.7548, 2013.

[7] V.J.W. Guo and C. Krattenthaler, *Some divisibility properties of binomial and q-binomial coefficients,* J. Number Theory 135 (2014), 167-184.

[8] V.J.W. Guo, F. Jouhet and J. Zeng, *Factors of alternating sums of products of binomial and q-binomial coefficients,* Acta Arith. 127 (2007), 17-31.

[9] V.J.W. Guo and J. Zeng, *Factors of binomial sums from the Catalan triangle,* J. Number Theory 130 (2010), 172-186.

[10] V.J.W. Guo and J. Zeng, *Factors of sums and alternating sums involving binomial coefficients and powers of integers,* Int. Number Theory 7 (2011), 1959-1976.

[11] M. Razpet, *On divisibility of binomial coefficients,* Discrete Math. 135 (1994), 377-379.

[12] Z.-W. Sun, *On divisibility of binomial coefficients,* J. Austral. Math. Soc. 93 (2012), 189-201.

[13] Z.-W. Sun, *Products and sums divisible by central binomial coefficients,* Electron. J. Combin. 20(1) (2013), #P9.